


**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

75-091

APPROVED DRAFT LABELING

CBLVER:R5




699

**CARBIDOPA and
LEVODOPA**
**Extended-release
Tablets**
500 mg/200 mg

Rx only


DESCRIPTION: Carbidopa and levodopa extended-release tablets are for the treatment of Parkinson's disease and syndrome.

Carbidopa, an inhibitor of aromatic amino acid decarboxylation, is a white, crystalline compound, slightly soluble in water, with a molecular weight of 244.25. It is designated chemically as (S)-1-(3,4-dihydroxyphenyl)-2-methyl-2-(propanoic acid monohydrate) propanoic acid monohydrate. Its structural formula is:



Tablet content is expressed in terms of anhydrous carbidopa, which has a molecular weight of 226.23.

Levodopa, an aromatic amino acid, is a white, crystalline compound, slightly soluble in water, with a molecular weight of 197.19. It is designated chemically as (S)-1-(3,4-dihydroxyphenyl)-2-(amino-3,4-dihydroxybenzyl) propanoic acid. Its structural formula is:



Each extended-release tablet, for oral administration, contains 500 mg of carbidopa and 200 mg of levodopa. In addition, each tablet contains the following inactive ingredients: FD&C Blue #2 Aluminum Lake, FD&C Red #40 Aluminum Lake, hydroxypropyl cellulose, hydroxypropyl methylcellulose, and magnesium stearate.

USP Drug release test pending.
CLINICAL PHARMACOLOGY: Mechanism of Action: Parkinson's disease is a progressive, neurodegenerative disorder of the extrapyramidal nervous system affecting the mobility and control of the skeletal muscular system. Its characteristic features include resting tremor, rigidity, and bradykinetic movements. Symptomatic treatments, such as levodopa therapies, may permit the patient better mobility.

Current evidence indicates that symptoms of Parkinson's disease are related to depletion of dopamine in the corpus striatum. Administration of dopamine is ineffective in the treatment of Parkinson's disease apparently because it does not cross the blood-brain barrier. However, levodopa, the metabolic precursor of dopamine, does cross the blood-brain barrier, and presumably is converted to dopamine in the brain. This is thought to be the mechanism whereby levodopa relieves symptoms of Parkinson's disease.

Pharmacodynamics: When levodopa is administered orally it is rapidly decarboxylated to dopamine in extracerebral tissues so that only a small portion of a given dose is transported unchanged to the central nervous system. For this reason, large doses of levodopa are required for adequate therapeutic effect and these may often be accompanied by nausea and other adverse reactions, some of which are attributable to dopamine formed in extracerebral tissues.

Since levodopa competes with certain amino acids for transport across the gut wall, the absorption of levodopa may be impaired in some patients on a high protein diet.

Carbidopa inhibits decarboxylation of peripheral levodopa. It does not cross the

live in the treatment of Parkinson's disease apparently because it does not cross the blood-brain barrier. However, levodopa, the metabolic precursor of dopamine, does cross the blood-brain barrier, and presumably is converted to dopamine in the brain. This is thought to be the mechanism whereby levodopa relieves symptoms of Parkinson's disease.

Pharmacodynamics: When levodopa is administered orally it is rapidly decarboxylated to dopamine in extracerebral tissues so that only a small portion of a given dose is transported unchanged to the central nervous system. For this reason, large doses of levodopa are required for adequate therapeutic effect and these may often be accompanied by nausea and other adverse reactions, some of which are attributable to dopamine formed in extracerebral tissues.

Since levodopa competes with certain amino acids for transport across the gut wall, the absorption of levodopa may be impaired in some patients on a high protein diet.

Carbidopa inhibits decarboxylation of peripheral levodopa. It does not cross the blood-brain barrier and does not affect the metabolism of levodopa within the central nervous system.

Since its decarboxylase-inhibiting activity is limited to extracerebral tissues, administration of carbidopa with levodopa makes more levodopa available for transport to the brain.

Patients treated with levodopa therapy for Parkinson's disease may develop motor fluctuations characterized by end-of-dose failure, peak dose dyskinesia, and akinesia. The advanced form of motor fluctuations ("on-off" phenomenon) is characterized by unpredictable swings from mobility to immobility. Although the causes of the motor fluctuations are not completely understood, in some patients they may be attenuated by treatment regimens that produce steady plasma levels of levodopa.

Each tablet contains 50 mg of carbidopa and 200 mg of levodopa, in a sustained-release dosage form designed to release these ingredients over a 4 to 6 hour period. With carbidopa and levodopa extended-release there is less variation in plasma levodopa levels than with carbidopa and levodopa immediate-release, the conventional formulation. However, carbidopa and levodopa extended-release is less systemically bioavailable than carbidopa and levodopa immediate-release and may require increased daily doses to achieve the same level of symptomatic relief as provided by carbidopa and levodopa immediate-release.

In clinical trials, patients with moderate to severe motor fluctuations who received carbidopa and levodopa extended-release did not experience quantitatively significant reductions in "off" time when compared to carbidopa and levodopa immediate-release. However, global ratings of improvement as assessed by both patient and physician were better during therapy with carbidopa and levodopa extended-release than with carbidopa and levodopa immediate-release. In patients without motor fluctuations, carbidopa and levodopa extended-release under controlled conditions, provided the same therapeutic benefit with less frequent dosing when compared to carbidopa and levodopa immediate-release.

Pharmacokinetics: Carbidopa reduces the amount of levodopa required to produce a given response by about 75 percent and, when administered with levodopa, increases both plasma levels and the plasma half-life of levodopa, and decreases plasma and urinary dopamine and homovanillic acid.

Elimination half-life of levodopa in the presence of carbidopa is about 1.5 hours. Following carbidopa and levodopa extended-release, the apparent half-life of levodopa may be prolonged because of continuous absorption.

In healthy elderly subjects (56 to 67 years old) the mean time-to-peak concentration of levodopa after a single dose of 50 mg/200 mg carbidopa and levodopa extended-release was about 2 hours as compared to 0.5 hours after standard carbidopa and levodopa immediate-release. The maximum concentration of levodopa after a single dose of carbidopa and levodopa extended-release was about 35% of the standard carbidopa and levodopa (1151 vs 3256 ng/mL). The extent of availability of levodopa from carbidopa and levodopa extended-release was about 70 to 75% relative to intravenous levodopa or standard carbidopa and levodopa immediate-release in the elderly. The absolute bioavailability of levodopa from carbidopa and levodopa extended-release (relative to IV) in young subjects was shown to be only about 44%. The extent of availability and the peak concentrations of levodopa were comparable in the elderly after a single dose and at steady state after L.d. administration of 50 mg/200 mg carbidopa and levodopa extended-release. In elderly subjects, the average trough levels of levodopa at steady state after the extended-release tablet were about 2 fold higher than after the standard carbidopa and levodopa immediate-release (153 vs 74 ng/mL).

In these studies, using similar total daily doses of levodopa, plasma levodopa concentrations with carbidopa and levodopa extended-release fluctuated in a narrower range than with carbidopa and levodopa immediate-release. Because the bioavailability of levodopa from carbidopa and levodopa extended-release relative to carbidopa and levodopa immediate-release is approximately 70 to 75%, the daily dosage of levodopa necessary to produce a given clinical response with the extended-release formulation will usually be higher.

The extent of availability and peak concentrations of levodopa after a single dose of 50 mg/200 mg carbidopa and levodopa extended-release increased by about 50% and 25%, respectively, when administered with food.

At steady state, the bioavailability of carbidopa from carbidopa and levodopa immediate-release is approximately 99% relative to the concomitant administration of carbidopa and levodopa. At steady state, carbidopa bioavailability from carbidopa and levodopa extended-release 50 mg/200 mg is

of levodopa from carbidopa and levodopa extended-release relative to carbidopa and levodopa immediate-release is approximately 70 to 75%, the daily dosage of levodopa necessary to produce a given clinical response with the extended-release formulation will usually be higher.

The extent of availability and peak concentrations of levodopa after a single dose of 50 mg/200 mg carbidopa and levodopa extended-release increased by about 50% and 75%, respectively, when administered with food.

At steady state, the bioavailability of carbidopa from carbidopa and levodopa immediate-release is approximately 95% relative to the concomitant administration of carbidopa and levodopa. At steady state, carbidopa bioavailability from carbidopa and levodopa extended-release 50 mg/200 mg is approximately 58% relative to that from carbidopa and levodopa immediate-release.

Pyridoxine hydrochloride (vitamin B₆), in oral doses of 10 mg to 25 mg, may reverse the effects of levodopa by increasing the rate of automatic amino acid decarboxylation. Carbidopa inhibits this action of pyridoxine.

INDICATIONS AND USAGE: Carbidopa and levodopa extended-release tablets are indicated in the treatment of the symptoms of idiopathic Parkinson's disease (paralysis agitans), postencephalitic parkinsonism, and symptomatic parkinsonism which may follow injury to the nervous system by carbon monoxide intoxication and/or manganese intoxication.

CONTRAINDICATIONS: Nonselective MAO inhibitors are contraindicated for use with carbidopa and levodopa extended-release. These inhibitors must be discontinued at least two weeks prior to initiating therapy with carbidopa and levodopa extended-release. Carbidopa and levodopa extended-release may be administered concomitantly with the manufacturer's recommended dose of an MAO inhibitor with selectivity for MAO type B (e.g., selegiline HCl) (see PRECAUTIONS Drug Interactions).

Carbidopa and levodopa extended-release is contraindicated in patients with known hypersensitivity to any component of this drug and in patients with narrow angle glaucoma.

Because levodopa may activate a malignant melanoma, carbidopa and levodopa extended-release should not be used in patients with suspicious, undiagnosed skin lesions or a history of melanoma.

WARNINGS: When patients are receiving levodopa without a decarboxylase inhibitor, levodopa must be discontinued at least twelve hours before carbidopa and levodopa extended-release is started. In order to reduce adverse reactions, it is necessary to individualize therapy. Carbidopa and levodopa extended-release should be substituted at a dosage that will provide approximately 75 percent of the previous levodopa dosage (see DOSAGE AND ADMINISTRATION).

Carbidopa does not decrease adverse reactions due to central effects of levodopa. By permitting more levodopa to reach the brain, particularly when nausea and vomiting is not a dose-limiting factor, certain adverse CNS effects, e.g., dyskinesias, will occur at lower dosages and sooner during therapy with carbidopa and levodopa extended-release than with levodopa alone.

Patients receiving carbidopa and levodopa extended-release may develop increased dyskinesias compared to carbidopa and levodopa immediate-release. Dyskinesias are a common side effect of carbidopa-levodopa treatment. The occurrence of dyskinesias may require dosage reduction.

As with levodopa, carbidopa and levodopa extended-release may cause mental disturbances. These reactions are thought to be due to increased brain dopamine following administration of levodopa. All patients should be observed carefully for the development of depression with concomitant suicidal tendencies. Patients with past or current psychoses should be treated with caution.

Carbidopa and levodopa extended-release should be administered cautiously to patients with severe cardiovascular or pulmonary disease, bronchial asthma, renal, hepatic or endocrine disease.

As with levodopa, care should be exercised in administering carbidopa and levodopa extended-release to patients with a history of myocardial infarction who have residual atrial, nodal, or ventricular arrhythmias. In such patients, cardiac function should be monitored with particular care during the period of initial dosage adjustment, in a facility with provisions for intensive cardiac care.

As with levodopa, treatment with carbidopa and levodopa extended-release may increase the possibility of upper gastrointestinal hemorrhage in patients with a history of peptic ulcer.

Neuroleptic Malignant Syndrome (NMS): Sporadic cases of a symptom complex resembling NMS have been reported in association with dose reductions or withdrawal of carbidopa and levodopa immediate-release and carbidopa and levodopa extended-release.

Therefore, patients should be observed carefully when the dosage of carbidopa and levodopa extended-release is reduced abruptly or discontinued, especially if the patient is receiving neuroleptics.

NMS is an uncommon but life-threatening syndrome characterized by fever or hyperthermia. Neurological findings, including muscle rigidity, involuntary movements, altered consciousness, mental status changes, other disturbances, such as autonomic dysfunction, tachycardia, tachypnea, sweating, hyper- or hypotension; laboratory findings, such as creatine phosphokinase elevation, leukocytosis, myoglobinuria, and increased serum myoglobin have been reported.

The early diagnosis of this condition is important for the appropriate management of these patients. Considering NMS as a possible diagnosis and ruling out other acute illnesses (e.g., pneumonia, systemic infection, etc.) is essential. This may be especially complete if the clinical presentation includes both serious medical illness and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat

4
mal, choreoathetoid, involuntary movements, rigidity, involuntary movements, ataxic consciousness, mental status changes, other disturbances, such as autonomic dysfunction, tachycardia, tachypnea, sweating, hyper- or hypotension, laboratory findings, such as creatine phosphokinase elevation, leukocytosis, myoglobinuria, and increased serum myoglobin have been reported.

The early diagnosis of this condition is important for the appropriate management of these patients. Considering NMS as a possible diagnosis and ruling out other acute illnesses (e.g., pneumonia, systemic infection, etc.) is essential. This may be especially complex if the clinical presentation includes both serious medical illness and untreated or inadequately treated extrapyramidal signs and symptoms (EPS). Other important considerations in the differential diagnosis include central anticholinergic toxicity, heat stroke, drug fever, and primary central nervous system (CNS) pathology.

The management of NMS should include: 1) intensive symptomatic treatment and medical monitoring and 2) treatment of any concomitant serious medical problems for which specific treatments are available. Dopamine agonists, such as bromocriptine, and muscle relaxants, such as dantrolene, are often used in the treatment of NMS, however, their effectiveness has not been demonstrated in controlled studies.

PRECAUTIONS: General: As with levodopa, periodic evaluations of hepatic, hematopoietic, cardiovascular, and renal function are recommended during extended therapy.

Patients with chronic wide angle glaucoma may be treated cautiously with carbidopa and levodopa provided the intraocular pressure is well controlled and the patient is monitored carefully for changes in intraocular pressure during therapy.

Information for Patients: The patient should be informed that carbidopa and levodopa extended-release tablets are a sustained-release formulation of carbidopa and levodopa which releases these ingredients over a 4 to 6 hour period. It is important that carbidopa and levodopa extended-release be taken at regular intervals according to the schedule outlined by the physician. The patient should be cautioned not to change the prescribed dosage regimen and not to add any additional antiparkinson medications, including other carbidopa and levodopa preparations, without first consulting the physician.

If abnormal involuntary movements appear or get worse during treatment with carbidopa and levodopa extended-release, the physician should be notified, as dosage adjustment may be necessary.

Patients should be advised that sometimes the onset of effect of the first morning dose of carbidopa and levodopa extended-release may be delayed for up to 1 hour compared with the response usually obtained from the first morning dose of carbidopa and levodopa immediate-release. The physician should be notified if such delayed responses pose a problem in treatment.

Patients should be advised that, occasionally, dark color (red, brown, or black) may appear in saliva, urine, or sweat after ingestion of carbidopa and levodopa extended-release. Although the color appears to be clinically insignificant, garments may become discolored.

The patient should be informed that a change in diet to foods that are high in protein may delay the absorption of levodopa and may reduce the amount taken up in the circulation. Excessive acidity also delays stomach emptying, thus delaying the absorption of levodopa. Iron salts (such as in multi-vitamin tablets) may also reduce the amount of levodopa available to the body. The above factors may reduce the clinical effectiveness of the levodopa or carbidopa-levodopa therapy.

Patients must be advised that the whole or half tablet should be swallowed without chewing or crushing.

NOTE: The suggested advice to patients being treated with carbidopa and levodopa extended-release tablets is intended to aid in the safe and effective use of this medication. It is not a disclosure of all possible adverse or intended effects.

Laboratory Tests: Abnormalities in laboratory tests may include elevations of liver function tests such as alkaline phosphatase, SGOT (AST), SGPT (ALT), lactic dehydrogenase, and bilirubin. Abnormalities in blood urea nitrogen and positive Coombs test have also been reported. Commonly, levels of blood urea nitrogen, creatinine, and uric acid are lower during administration of carbidopa and levodopa preparations than with levodopa.

Carbidopa and levodopa preparations may cause a false-positive reaction for urinary ketone bodies when a test tape is used for determination of ketonuria. This reaction will not be altered by boiling the urine specimen. False-negative tests may result with the use of glucose-oxidase methods of testing for glucosuria.

Cases of falsely diagnosed pheochromocytoma in patients on carbidopa-levodopa therapy have been reported very rarely. Caution should be exercised when interpreting the plasma and urine levels of catecholamines and their metabolites in patients on levodopa or carbidopa-levodopa therapy.

Drug Interactions: Caution should be exercised when the following drugs are administered concomitantly with carbidopa and levodopa.

Symptomatic postural hypotension has occurred when carbidopa and levodopa preparations were added to the treatment of patients receiving some antihypertensive drugs. Therefore, when therapy with carbidopa and levodopa is started, dosage adjustment of the antihypertensive drug may be required.

For patients receiving monoamine oxidase (MAO) inhibitors (Type A or B), see **CONTRAINDICATIONS**. Concomitant therapy with selegiline and carbidopa-levodopa may be associated with severe orthostatic hypotension not attributable to carbidopa-levodopa alone (see **CONTRAINDICATIONS**).

There have been rare reports of adverse reactions, including hypertension and dyskinesia, resulting from the concomitant use of tricyclic antidepressants and carbidopa and levodopa preparations.

5

observed in controlled studies.

PRECAUTIONS: General. As with levodopa, periodic evaluations of hepatic, hematopoietic, cardiovascular, and renal function are recommended during extended therapy.

Patients with chronic wide angle glaucoma may be treated cautiously with carbidopa and levodopa provided the intraocular pressure is well controlled and the patient is monitored carefully for changes in intraocular pressure during therapy.

Information for Patients: The patient should be informed that carbidopa and levodopa extended-release tablets are a sustained-release formulation of carbidopa and levodopa which releases these ingredients over a 4 to 8 hour period. It is important that carbidopa and levodopa extended-release be taken at regular intervals according to the schedule outlined by the physician. The patient should be cautioned not to change the prescribed dosage regimen and not to add any additional antiparkinsonian medications, including other carbidopa and levodopa preparations, without first consulting the physician.

If abnormal involuntary movements appear or get worse during treatment with carbidopa and levodopa extended-release, the physician should be notified, as dosage adjustment may be necessary.

Patients should be advised that sometimes the onset of effect of the first morning dose of carbidopa and levodopa extended-release may be delayed for up to 1 hour compared with the response usually obtained from the first morning dose of carbidopa and levodopa immediate-release. The physician should be notified if such delayed responses pose a problem in treatment.

Patients should be advised that, occasionally, dark color (red, brown, or black) may appear in saliva, urine, or sweat after ingestion of carbidopa and levodopa extended-release. Although the color appears to be clinically insignificant, garments may become discolored.

The patient should be informed that a change in diet to foods that are high in protein may delay the absorption of levodopa and may reduce the amount taken up in the circulation. Excessive acidity also delays stomach emptying, thus delaying the absorption of levodopa. Iron salts (such as in multi-vitamin tablets) may also reduce the amount of levodopa available to the body. The above factors may reduce the clinical effectiveness of the levodopa or carbidopa-levodopa therapy.

Patients must be advised that the whole or half tablet should be swallowed without chewing or crushing.

NOTE: The suggested advice to patients being treated with carbidopa and levodopa extended-release tablets is intended to aid in the safe and effective use of this medication. It is not a disclosure of all possible adverse or intended effects.

Laboratory Tests: Abnormalities in laboratory tests may include elevations of liver function tests such as alkaline phosphatase, SGOT (AST), SGPT (ALT), lactic dehydrogenase, and bilirubin. Abnormalities in blood urea nitrogen and positive Coombs test have also been reported. Commonly, levels of blood urea nitrogen, creatinine, and uric acid are lower during administration of carbidopa and levodopa preparations than with levodopa.

Carbidopa and levodopa preparations may cause a false-positive reaction for urinary ketone bodies when a test tape is used for determination of ketonuria. This reaction will not be altered by boiling the urine specimen. False-negative tests may result with the use of glucose-oxidase methods of testing for glucosuria.

Cases of falsely diagnosed pheochromocytoma in patients on carbidopa-levodopa therapy have been reported very rarely. Caution should be exercised when interpreting the plasma and urine levels of catecholamines and their metabolites in patients on levodopa or carbidopa-levodopa therapy.

Drug Interactions: Caution should be exercised when the following drugs are administered concomitantly with carbidopa and levodopa.

Symptomatic postural hypotension has occurred when carbidopa and levodopa preparations were added to the treatment of patients receiving some antihypertensive drugs. Therefore, when therapy with carbidopa and levodopa is started, dosage adjustment of the antihypertensive drug may be required.

For patients receiving monoamine oxidase (MAO) inhibitors (Type A or B), see **CONTRAINDICATIONS**. Concomitant therapy with selegiline and carbidopa-levodopa may be associated with severe orthostatic hypotension not attributable to carbidopa-levodopa alone (see **CONTRAINDICATIONS**).

There have been rare reports of adverse reactions, including hypertension and dyskinesia, resulting from the concomitant use of tricyclic antidepressants and carbidopa and levodopa preparations.

Dopamine D₂ receptor antagonists (e.g., phenothiazines, butyrophenones, risperidone) and isoniazid may reduce the therapeutic effects of levodopa. In addition, the beneficial effects of levodopa in Parkinson's disease have been reported to be reversed by phenothiazine and papaverine. Patients taking these drugs with carbidopa and levodopa should be carefully observed for loss of therapeutic response.

Iron salts may reduce the bioavailability of levodopa and carbidopa. The clinical relevance is unclear.

Although metoclopramide may increase the bioavailability of levodopa by increasing gastric emptying, metoclopramide may also

adversely affect disease control by its dopamine receptor antagonistic properties. **Carcinogenesis, Mutagenesis, Impairment of Fertility:** In a two-year bioassay of carbidopa and levodopa immediate-release, no evidence of carcinogenicity was found in rats receiving doses of approximately two times the maximum daily human dose of carbidopa and four times the maximum daily human dose of levodopa (equivalent to 8 carbidopa and levodopa extended-release tablets).

In reproduction studies with carbidopa and levodopa immediate-release, no effects on fertility were found in rats receiving doses of approximately two times the maximum daily human dose of carbidopa and four times the maximum daily human dose of levodopa (8 carbidopa and levodopa extended-release tablets).

Pregnancy: Teratogenic Effects. Pregnancy Category C: No teratogenic effects were observed in a study in mice receiving up to 20 times the maximum recommended human dose of carbidopa and levodopa immediate-release. There was a decrease in the number of live pups delivered by rats receiving approximately two times the maximum recommended human dose of carbidopa and approximately five times the maximum recommended human dose of levodopa during organogenesis. Carbidopa and levodopa immediate-release caused both visceral and skeletal malformations in rabbits at all doses and ratios of carbidopa/levodopa tested, which ranged from 10 times/5 times the maximum recommended human dose of carbidopa/levodopa to 20 times/10 times the maximum recommended human dose of carbidopa/levodopa.

There are no adequate or well-controlled studies in pregnant women. It has been reported from individual cases that levodopa crosses the human placental barrier, enters the fetus, and is metabolized. Carbidopa concentrations in fetal tissue appeared to be minimal. Use of carbidopa and levodopa extended-release in women of childbearing potential requires that the anticipated benefits of the drug be weighed against possible hazards to mother and child.

Nursing Mothers: It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when carbidopa and levodopa extended-release is administered to a nursing woman.

Pediatric Use: Safety and effectiveness in pediatric patients have not been established. Use of the drug in patients below the age of 18 is not recommended.

ADVERSE REACTIONS: In controlled clinical trials, patients predominantly with moderate to severe motor fluctuations while on carbidopa and levodopa immediate-release were randomized to therapy with either carbidopa and levodopa immediate-release or carbidopa and levodopa extended-release. The adverse experience frequency profile of carbidopa and levodopa extended-release did not differ substantially from that of carbidopa and levodopa immediate release, as shown in Table I.

Table I
Clinical Adverse Experiences Occurring in 1% or Greater of Patients

Adverse Experience	Carbidopa and Levodopa Extended-release n=491 %	Carbidopa and Levodopa Immediate-release n=524 %
Dyskinesia	16.5	12.2
Nausea	5.5	5.7
Hallucinations	3.9	3.2
Confusion	3.7	2.3
Dizziness	2.9	2.3
Depression	2.2	1.3
Urinary tract infection	2.2	2.3
Headache	2.0	1.9
Dream abnormalities	1.8	0.8
Dystonia	1.8	0.8
Vomiting	1.8	1.9
Upper respiratory infection	1.8	1.0
Dyspnea	1.6	0.4
"On-Off" phenomena	1.6	1.1
Back pain	1.6	0.6
Dry mouth	1.4	1.1
Anorexia	1.2	1.1
Diarrhea	1.2	0.6
Insomnia	1.2	1.0
Orthostatic hypotension	1.0	1.1
Shoulder pain	1.0	0.6
Chest pain	1.0	0.8
Muscle cramps	0.8	1.0
Paresthesia	0.8	1.1
Urinary frequency	0.8	1.1
Dyspepsia	0.6	1.1
Constipation	0.2	1.5

1. Abnormal laboratory findings occurring at a frequency of 1% or greater in approximately 443 patients who received carbidopa and levodopa extended-release and 475 who received carbidopa and levodopa immediate-release during controlled clinical trials included: decreased hemoglobin and hematocrit; elevated serum glucose; white blood cells; bacteria and blood in the urine.

2. The adverse experiences observed in patients in uncontrolled studies were similar to those seen in controlled clinical studies.

Other adverse experiences reported overall in clinical trials in 748 patients treated with carbidopa and levodopa extended-release, listed by body system in order of decreasing frequency, include:

Body as a Whole: Asthenia, fatigue, abdominal pain, orthostatic effects.
Cardiovascular: Palpitation, hypertension, hypotension, myocardial infarction.
Gastrointestinal: Gastrointestinal pain, dysphagia, heartburn.
Metabolic: Weight loss.
Musculoskeletal: Leg pain

Adverse	1.1	1.1
Diarrhea	1.2	0.6
Insomnia	1.2	1.0
Orthostatic		
hypertension	1.0	1.1
Shoulder pain	1.0	0.6
Chest pain	1.0	0.8
Muscle cramps	0.8	1.0
Paresthesia	0.8	1.1
Urinary frequency	0.8	1.1
Dyspepsia	0.6	1.1
Constipation	0.2	1.5

1. Abnormal laboratory findings occurring at a frequency of 1% or greater in approximately 443 patients who received carbidopa and levodopa extended-release and 475 who received carbidopa and levodopa immediate-release during controlled clinical trials included: decreased hemoglobin and hematocrit; elevated serum glucose, white blood cells, bacteria and blood in the urine.

2. The adverse experiences observed in patients in uncontrolled studies were similar to those seen in controlled clinical studies.

Other adverse experiences reported overall in clinical trials in 748 patients treated with carbidopa and levodopa extended-release, listed by body system in order of decreasing frequency, include:

Body as a Whole: Asthenia, fatigue, abdominal pain, orthostatic effects.

Cardiovascular: Palpitation, hypertension, hypotension, myocardial infarction.

Gastrointestinal: Gastrointestinal pain, dysphagia, heartburn.

Metabolic: Weight loss.

Musculoskeletal: Leg pain.

Nervous System/Psychiatric: Chorea, somnolence, falling, anxiety, disorientation, decreased mental acuity, gait abnormalities, encephalomyelitis, agitation, nervousness, sleep disorders, memory impairment.

Respiratory: Cough, pharyngeal pain, common cold.

Skin: Rash.

Special Senses: Blurred vision.

Urogenital: Urinary incontinence.

Laboratory tests: Decreased white blood cell count and serum potassium; increased BUN, serum creatinine and serum LDH; protein and glucose in the urine.

The following adverse experiences have been reported in post-marketing experience with carbidopa and levodopa extended-release:

Cardiovascular: Cardiac irregularities, syncope.

Gastrointestinal: Taste alterations, dark saliva.

Hypersensitivity: Angioedema, urticaria, pruritus.

Nervous System/Psychiatric: Neuroleptic malignant syndrome (see WARNINGS), increased tremor, peripheral neuropathy, psychotic episodes including delusions and paranoid ideation.

Skin: Alopecia, flushing, dark sweat.

Urogenital: Dark urine.

Other adverse reactions that have been reported with levodopa alone and with various carbidopa-levodopa formulations and may occur with carbidopa and levodopa extended-release are:

Cardiovascular: Phlebitis.

Gastrointestinal: Gastrointestinal bleeding, development of duodenal ulcer, sialorrhea, bruxism, hiccups, flatulence, burning sensation of tongue.

Hematologic: Hemolytic and nonhemolytic anemia, thrombocytopenia, leukopenia, agranulocytosis.

Hypersensitivity: Henoch-Schönlein purpura.

Metabolic: Weight gain, edema.

Nervous System/Psychiatric: Ataxia, depression with suicidal tendencies, dementia, euphoria, convulsions (however, a causal relationship has not been established); bradykinetic episodes, numbness, muscle twitching, blepharospasm (which may be taken as an early sign of excess dosage; consideration of dosage reduction may be made at this time), trismus, activation of latent Horner's syndrome, nystagmus.

Skin: Malignant melanoma (see also CONTRAINDICATIONS), increased sweating.

Special Senses: Oculogyric crisis, mydriasis, diplopia.

Urogenital: Urinary retention, priapism.

Miscellaneous: Faintness, hoarseness, malaise, hot flashes, sense of stimulation, bizarre breathing patterns.

Laboratory tests: Abnormalities in alkaline phosphatase, SGOT (AST), SGPT (ALT), bilirubin, Coombs test, uric acid.

OVERDOSAGE: Management of acute overdosage with carbidopa and levodopa extended-release is the same as with levodopa. Pyridoxine is not effective in reversing the actions of carbidopa and levodopa extended-release.

General supportive measures should be employed, along with immediate gastric lavage. Intravenous fluids should be administered judiciously and an adequate airway maintained. Electrocardiographic monitoring should be instituted and the patient carefully observed for the development of arrhythmias; if required, appropriate antiarrhythmic therapy should be given. The possibility that the patient may have taken other drugs as well as carbidopa and levodopa extended-release should be taken into consideration. To date, no experience has been reported with dialysis; hence, its value in overdosage is not known.

Based on studies in which high doses of levodopa and/or carbidopa were administered, a significant proportion of rats and mice given single oral doses of levodopa of approximately 1500 to 2000 mg/kg are expected to die. A significant proportion of infant rats of both sexes are expected to die at a dose of 800 mg/kg. A significant proportion of rats are expected to die after treatment with similar doses of carbidopa. The addition of carbidopa in a 1:10 ratio with levodopa increases the dose at which a significant proportion of mice are expected to die to 3350 mg/kg.

DOSEAGE AND ADMINISTRATION: Carbidopa and levodopa extended-release tablets contain carbidopa (50 mg) and levodopa (200 mg) in a 1:4 ratio. The daily dosage of carbidopa and levodopa extended-release must be determined by careful titration. Patients should be monitored closely during

...to die. A significant proportion of infant rats of both sexes are expected to die at a dose of 800 mg/kg. A significant proportion of rats are expected to die after treatment with similar doses of carbidopa. The addition of levodopa in a 1:10 ratio with levodopa increases the dose at which a significant proportion of mice are expected to die to 3360 mg/kg.

DOSEAGE AND ADMINISTRATION: Carbidopa and levodopa extended-release tablets contain carbidopa (50 mg) and levodopa (500 mg) in a 1:10 ratio. The daily dosage of carbidopa and levodopa extended-release must be determined by careful titration. Patients should be monitored closely during the dose adjustment period, particularly with regard to appearance or worsening of involuntary movements, dyskinesias or nausea. Carbidopa and levodopa extended-release tablets may be administered as whole or as half-tablets which should not be chewed or crushed.

Standard drugs for Parkinson's disease, other than levodopa without a decarboxylase inhibitor, may be used concomitantly while carbidopa and levodopa extended-release is being administered, although their dosage may have to be adjusted.

Since carbidopa prevents the reversal of levodopa effects caused by pyridoxine, carbidopa and levodopa extended-release can be given to patients receiving supplemental pyridoxine (vitamin B₆).

Initial Dosage: Patients Currently Treated with Conventional Carbidopa-Levodopa Preparations: Studies show that peripheral dopa-decarboxylase is saturated by the bioavailable carbidopa at doses of 70 mg a day and greater. Because the bioavailabilities of carbidopa and levodopa in carbidopa and levodopa immediate-release and carbidopa and levodopa extended-release are different, appropriate adjustments should be made, as shown in Table II.

Label	Approximate Bioavailability at Steady State*	
	Amount of Levodopa (mg) in Each Tablet	Approximate Amount of Bioavailable Levodopa (mg) in Each Tablet
Carbidopa and Levodopa Extended-release 50 mg/500 mg	200	100
Carbidopa and Levodopa Immediate-release 25 mg/250 mg	100	99***

* This table is only a guide to bioavailabilities since other factors such as food, drugs, and inter-patient variables may affect the bioavailability of carbidopa and levodopa.

** The extent of availability of levodopa from carbidopa and levodopa extended-release was about 70 to 75% relative to intravenous levodopa or standard carbidopa and levodopa immediate-release in the elderly.

*** The extent of availability of levodopa from carbidopa and levodopa immediate-release was 99% relative to intravenous levodopa in the healthy elderly.

Dosage with carbidopa and levodopa extended-release should be substituted at an amount that provides approximately 10% more levodopa per day, although this may need to be increased to a dosage that provides up to 30% more levodopa per day depending on clinical response (see DOSEAGE AND ADMINISTRATION: Titration). The interval between doses of carbidopa and levodopa extended-release should be 4 to 8 hours during the waking day. (See CLINICAL PHARMACOLOGY: Pharmacodynamics.)

A guideline for titration of carbidopa and levodopa extended-release is shown in Table III.

Table III
Guidelines for Initial Conversion From Carbidopa and Levodopa Immediate-release to Carbidopa and Levodopa Extended-release

Carbidopa and Levodopa Immediate-release Total Daily Dose* (mg)	Carbidopa and Levodopa Extended-release Suggested Dosage Regimen
300-400	1 tab b.i.d.
500-600	1 1/2 tab b.i.d. or 2 tab t.i.d.
700-800	A total of 4 tabs in 3 or more divided doses (e.g., 1 1/2 tab a.m., 1 1/2 tab early p.m., and 1 tab later p.m.)
900-1000	A total of 5 tabs in 3 or more divided doses (e.g., 2 tabs a.m., 2 tabs early p.m., and 1 tab later p.m.)

* For dosing ranges not shown in the table see DOSEAGE AND ADMINISTRATION: Initial Dosage: Patients Currently Treated with Conventional Carbidopa and Levodopa Preparations.

Patients Currently Treated with Levodopa Without a Decarboxylase Inhibitor: Levodopa must be discontinued at least twelve hours before therapy with carbidopa and levodopa extended-release is started. Carbidopa and levodopa extended-release should be substituted at a dosage that will provide approximately 25% of the previous levodopa dosage. In patients with mild to moderate disease, the initial dose is usually 1 tablet of carbidopa and levodopa extended-release tablets b.i.d.

9

doses (e.g., 1 1/2
tab a.m., 1 1/2 tab
early p.m., and
1 tab later p.m.)
A total of 3 tabs in 3
or more divided doses
(e.g., 2 tabs a.m.,
2 tabs early p.m., and
1 tab later p.m.)

* For dosing ranges not shown in the table
see DOSAGE AND ADMINISTRATION: Initial
Dosage. Patients Currently Treated with
Conventional Carbidopa and Levodopa
Preparations.

**Patients Currently Treated with Levodopa
Without a Decarboxylase Inhibitor:** Levo-
dopa must be discontinued at least twelve
hours before therapy with carbidopa and
levodopa extended-release is started.
Carbidopa and levodopa extended-release
should be substituted at a dosage that will
provide approximately 25% of the previous
levodopa dosage. In patients with mild to
moderate disease, the initial dose is usually
1 tablet of carbidopa and levodopa extend-
ed-release tablets b.i.d.

Patients Not Receiving Levodopa: In pa-
tients with mild to moderate disease, the ini-
tial recommended dose is 1 tablet of carbi-
dopa and levodopa extended-release tablets
b.i.d. Initial dosage should not be given at
intervals of less than 6 hours.

**Titration with Carbidopa and Levodopa
Extended-release:** Following initiation of
therapy, doses and dosing intervals may be
increased or decreased depending upon
therapeutic response. Most patients have
been adequately treated with doses of carbi-
dopa and levodopa extended-release that
provide 400 to 1600 mg of levodopa per day,
administered as divided doses at intervals
ranging from 4 to 8 hours during the waking
day. Higher doses of carbidopa and levodopa
extended-release (2400 mg or more of levo-
dopa per day) and shorter intervals (less
than 4 hours) have been used, but are not
usually recommended.

When doses of carbidopa and levodopa
extended-release are given at intervals of
less than 4 hours, and/or if the divided doses
are not equal, it is recommended that the
smaller doses be given at the end of the day.

An interval of at least 3 days between
dosage adjustments is recommended.

Maintenance: Because Parkinson's disease
is progressive, periodic clinical evaluations
are recommended; adjustment of the dose
regimen of carbidopa and levodopa
extended-release may be required.

**Addition of Other Antiparkinson Medi-
cations:** Anticholinergic agents, dopamine
agonists, and amantadine can be given with
carbidopa and levodopa extended-release.
Dosage adjustment of carbidopa and levo-
dopa extended-release may be necessary when
these agents are added.

A dose of carbidopa and levodopa imme-
diate-release 25-100 or 10-100 (one half or
a whole tablet) can be added to the dosage
regimen of carbidopa and levodopa extend-
ed-release in selected patients with ad-
vanced disease who need additional imme-
diate-release levodopa for a brief time dur-
ing daytime hours.

Interruption of Therapy: Sporadic cases of a
symptom complex resembling Neuroleptic
Malignant Syndrome (NMS) have been asso-
ciated with dose reductions and withdrawal
of carbidopa and levodopa immediate-
release or carbidopa and levodopa extended-
release.

Patients should be observed carefully if
abrupt reduction or discontinuation of carbi-
dopa and levodopa extended-release is
required, especially if the patient is receiving
neuroleptics. (See WARNINGS.)

If general anesthesia is required, carbi-
dopa and levodopa extended-release may be
continued as long as the patient is permitted
to take oral medication. If therapy is inter-
rupted temporarily, the patient should be
observed for symptoms resembling NMS, and
the usual dosage should be administered as
soon as the patient is able to take oral med-
ication.

HOW SUPPLIED: Carbidopa and Levodopa
Extended-release Tablets containing 50 mg
of carbidopa and 200 mg of levodopa are
available as purple oval, scored, biconvex
tablets debossed with MYLAN on one side of
the tablet and 9 to the left of the score and 4
to the right of the score on the other side of
the tablet. They are available as follows:

NDC 0378-0094-01
bottles of 100 tablets
NDC 0378-0094-05
bottles of 500 tablets

Dispense in a tight container as defined
in the USP using a child-resistance closure.
Keep container tightly closed.
DO NOT STORE ABOVE 30°C (86°F).



Mylan Pharmaceuticals Inc.
Morgantown, WV 26505

REVISED SEPTEMBER 1999
CBLVER.RS

orig
2

MYLAN PHARMACEUTICALS INC.


CARBIDOPA and LEVODOPA ER
TABLETS, 50/200mg

ANDA 75-091

Each extended-release
tablet contains:
Carbidopa 50 mg*
* (Anhydrous equivalent)
Levodopa 200 mg

50 mg/200 mg

N 3 0378-0094-05 5



NDC 0378-0094-05

MYLAN®

**CARBIDOPA and
LEVODOPA
EXTENDED-RELEASE
TABLETS
50 mg/200 mg**

500 TABLETS

CAUTION: Federal law prohibits
dispensing without prescription.

Dispense in a tight container as
defined in the USP using a child-
resistant closure.

Keep container tightly closed.

Keep this and all medication
out of the reach of children.

DO NOT STORE ABOVE 30°C (86°F).

Usual Adult Dosage: See accompanying
information.

Tablets should be swallowed without
chewing or crushing.


**Mylan Pharmaceuticals Inc.
Morgantown, WV 26505**

RM0094B

Each extended-release
tablet contains:
Carbidopa 50 mg*
* (Anhydrous equivalent)
Levodopa 200 mg

50 mg/200 mg

N 3 0378-0094-05 5



NDC 0378-0094-05

MYLAN®

**CARBIDOPA and
LEVODOPA
EXTENDED-RELEASE
TABLETS
50 mg/200 mg**

500 TABLETS

CAUTION: Federal law prohibits
dispensing without prescription.

Dispense in a tight container as
defined in the USP using a child-
resistant closure.

Keep container tightly closed.

Keep this and all medication
out of the reach of children.

DO NOT STORE ABOVE 30°C (86°F).

Usual Adult Dosage: See accompanying
information.

Tablets should be swallowed without
chewing or crushing.


**Mylan Pharmaceuticals Inc.
Morgantown, WV 26505**

RM0094B

Each extended-release
tablet contains:
Carbidopa 50 mg*
* (Anhydrous equivalent)
Levodopa 200 mg

50 mg/200 mg

N 3 0378-0094-05 5



NDC 0378-0094-05

MYLAN®

**CARBIDOPA and
LEVODOPA
EXTENDED-RELEASE
TABLETS
50 mg/200 mg**

500 TABLETS

CAUTION: Federal law prohibits
dispensing without prescription.

Dispense in a tight container as
defined in the USP using a child-
resistant closure.

Keep container tightly closed.

Keep this and all medication
out of the reach of children.

DO NOT STORE ABOVE 30°C (86°F).

Usual Adult Dosage: See accompanying
information.

Tablets should be swallowed without
chewing or crushing.

**Mylan Pharmaceuticals Inc.
Morgantown, WV 26505**

RM0094B

Mingo

MYLAN PHARMACEUTICALS INC.

CARBIDOPA and LEVODOPA ER
TABLETS, 50/200mg

ANDA 75-091

Each extended-release tablet contains:
Carbidopa 50 mg
(Anhydrous equivalent)
Levodopa 200 mg

50 mg/200 mg

SEP 30

NDC 0378-0094-01

MYLAN®

**CARBIDOPA and LEVODOPA
EXTENDED-RELEASE
TABLETS**
50 mg/200 mg

100 TABLETS

CAUTION: Federal law prohibits dispensing without prescription.
Dispense in a tight container as defined in the USP using a child-resistant closure.
Keep container tightly closed.
Keep this and all medication out of the reach of children.
DO NOT STORE ABOVE 30°C (86°F).
Usual Adult Dosage: See accompanying information.
Tablets should be swallowed without chewing or crushing.
Mylan Pharmaceuticals Inc.
Morgantown, WV 26505

RM0094A

Each extended-release tablet contains:
Carbidopa 50 mg
(Anhydrous equivalent)
Levodopa 200 mg

50 mg/200 mg

SEP 30

NDC 0378-0094-01

MYLAN®

**CARBIDOPA and LEVODOPA
EXTENDED-RELEASE
TABLETS**
50 mg/200 mg

100 TABLETS

CAUTION: Federal law prohibits dispensing without prescription.
Dispense in a tight container as defined in the USP using a child-resistant closure.
Keep container tightly closed.
Keep this and all medication out of the reach of children.
DO NOT STORE ABOVE 30°C (86°F).
Usual Adult Dosage: See accompanying information.
Tablets should be swallowed without chewing or crushing.
Mylan Pharmaceuticals Inc.
Morgantown, WV 26505

RM0094A

Each extended-release tablet contains:
Carbidopa 50 mg
(Anhydrous equivalent)
Levodopa 200 mg

50 mg/200 mg

SEP 30 199

NDC 0378-0094-01

MYLAN®

**CARBIDOPA and LEVODOPA
EXTENDED-RELEASE
TABLETS**
50 mg/200 mg

100 TABLETS

CAUTION: Federal law prohibits dispensing without prescription.
Dispense in a tight container as defined in the USP using a child-resistant closure.
Keep container tightly closed.
Keep this and all medication out of the reach of children.
DO NOT STORE ABOVE 30°C (86°F).
Usual Adult Dosage: See accompanying information.
Tablets should be swallowed without chewing or crushing.
Mylan Pharmaceuticals Inc.
Morgantown, WV 26505

RM0094A